

Cissoid *

History

Diocles (250 – ~100 BC) invented this curve to solve the doubling of the cube problem (also know as the the Delian problem). The name cissoid (ivy-shaped) derives from the shape of the curve. Later the method used to generate this curve was generalized, and we call all curves generated in a similar way cissoids. Newton (see below) found a way to generate the cissoid mechanically. The same kinematic motion with a different choice of drawing pin generates the (right) strophoid.

From Thomas L. Heath's Euclid's Elements translation (1925) (comments on definition 2, book one):

This curve is assumed to be the same as that by means of which, according to Eutocius, Diocles in his book On burning-glasses solved the problem of doubling the cube.

From Robert C. Yates' Curves and their properties (1952):

As early as 1689, J. C. Sturm, in his Mathesis Eucleata, gave a mechanical device for the constructions of the cissoid of Diocles.


From E.H.Lockwood A book of Curves (1961):

The name cissoid (“Ivy-shaped”) is mentioned by Geminus in the first century B.C., that is, about a century

*This file is from the 3D-XploreMath project.
Please see <http://rsp.math.brandeis.edu/3D-XplorMath/index.html>

after the death of the inventor Diocles. In the commentaries on the work by Archimedes *On the Sphere and the Cylinder*, the curve is referred to as Diocles' contribution to the classic problem of doubling the cube. ... Fermat and Roberval constructed the tangent (1634); Huygens and Wallis found the area (1658); while Newton gives it as an example, in his *Arithmetica Universalis*, of the ancients' attempts at solving cubic problems and again as a specimen in his *Enumeratio Linearum Tertii Ordinis*.

1 Description



The Cissoïd of Diocles is a special case of the general cissoïd. It is a cissoïd of a circle and a line tangent to the circle with respect to a point on the circle opposite to the tangent point. Here is a step-by-step description of the construction:

1. Let there be given a circle C and a line L tangent to this circle.
2. Let O be the point on the circle opposite to the tangent point.

3. Let $P1$ be a point on the circle C .
4. Let $P2$ be the intersection of line $[O,P1]$ and L .
5. Let Q be a point on line $[O,P1]$ such that

$$\text{dist}[O, Q] = \text{dist}[P1, P2].$$
6. The locus of Q (as $P1$ moves on C) is the cissoid of Diocles.

An important property to note is that Q and $P1$ are symmetric with respect to the midpoint of segment $[O,P2]$. Call this midpoint M . We can reflect every element in the construction around M , which will help us visually see other properties.

2 Formula derivation

Let the given circle C be centered at $(1/2, 0)$ with radius $1/2$. Let the given line L be $x = 1$, and let the given point O be the origin. Let $P1$ be a variable point on the circle, and Q the tracing point on line $[O, P1]$. Let the point $(1, 0)$ be A . We want to describe distance $r = \text{dist}[O, Q]$ in terms of the angle $\theta = [A, O, P1]$. This will give us an equation for the Cissoid in polar coordinates (r, θ) . From elementary geometry, the triangle $[A, O, P1]$ is a right triangle, so by trigonometry, the length of $[O, P1]$ is $\cos(\theta)$. Similarly, triangle $[O, A, P2]$ is a right triangle and the length of $[O, P2]$ is $\frac{1}{\cos(\theta)}$. Since $\text{dist}[O, Q] = \text{dist}[O, P2] - \text{dist}[O, P1]$, we have $\text{dist}[O, Q] = \frac{1}{\cos(\theta)} - \cos(\theta)$. Thus the polar equation is $r = \frac{1}{\cos(\theta)} - \cos(\theta)$. If we combine the fractions and use the identity $\sin^2 + \cos^2 = 1$, we arrive at an equivalent form: $r = \sin(\theta) \tan(\theta)$.

3 Formulas for Cissoid

In the following, the cusp is at the origin, and the asymptote is $x = 1$. (So the diameter of the circle is 1.)

Parametric: $(\sin^2(t), \sin^2(t) \tan(t)) \quad -\pi/2 < t < \pi/2$.

Parametric: $\left(\frac{t^2}{(1+t^2)}, \frac{t^3}{(1+t^2)}\right) \quad -\infty < t < \infty$

Strophoid: $\left(\frac{t^2-1}{(1+t^2)}, \frac{t(t^2-1)}{(1+t^2)}\right) \quad -\infty < t < \infty$

Polar: $r = \frac{1}{\cos(\theta)} - \cos(\theta) \quad -\pi/2 < t < \pi/2$.

Cisoid: $y^2(1-x) = x^3$, Strophoid: $y^2(1-x) = x^2(1+x)$.

The Cisoid has numerous interesting properties.

4 Properties

4.1 Doubling the Cube


Given a segment $[C, B]$, with the help of Cisoid of Diocles we can construct a segment $[C, M]$ such that $\text{dist}[C, M]^3 = 2 * \text{dist}[C, B]^3$. This solves the famous doubling the cube problem.

Step-by-step description:


1. Given two points C and B .
2. Construct a circle $c1$, centered on C and passing through B .
3. Construct points O and A on the circle such that line $[O, A]$ is perpendicular to line $[C, B]$

4. Construct a cissoid of Diocles using circle c_1 , tangent at A , and pole at O .
5. Construct point D such that B is the midpoint of segment $[C,D]$.
6. Construct line $[A,D]$. Let the intersection of cissoid and line $[A,D]$ be Q . (the intersection cannot be found with Greek Ruler and Compass. We assume it is a given.)
7. Let the intersection of line $[C, D]$ and line $[O, Q]$ be M .
8. $\text{dist}[C, M]^3 = 2 * \text{dist}[C, D]^3$

This can be proved trivially with analytic geometry.




4.2 Diocles' Construction



By some modern common accounts (Morris Kline, Thomas L. Heath), here's how Diocles constructed the curve in his book *On Burning-glasses*: Let AB and CD be perpendicular diameters of a circle. Let E be a point on arc $[B,C]$, and Z be a point on arc $[B,D]$, such that BE, BZ are equal. Draw ZH perpendicular to CD . Draw ED . Let P be intersection $[ZH,ED]$. The cissoid is the locus of all points P determined by all positions of E on arc $[B,C]$ and Z on arc $[B,D]$ with $\text{arc}[B,E]=\text{arc}[B,Z]$. (the portion of the curve that lies outside of the circle is a latter generalization).

In the curve, we have $CH/HZ=HZ/HD=HD/HP$. Thus HZ and HD are two mean proportionals between CH and HP . Proof: taking $CH/HZ=HZ/HD$, we have $CH * HD = HZ^2$. triangle $[D,C,Z]$ is a right triangle since it's a triangle on a circle with one side being the diameter (elementary geometry). We know an angle $[D,C,Z]$ and one side distance $[D,C]$, thus by trigonometry of right angles, we can derive all lengths DZ, CZ , and HZ . Substituting the results of computation in $CH * HD = HZ^2$ results an identity. Similarly, we know length HP and find $HZ/HD=HD/HP$ to be an identity.

4.3 Newton's Carpenter's Square and Tangent



Newton showed that Cissoid of Diocles and the right Strophoid can be generated by sliding a right triangle. The midpoint J of the edge CF draws the Cissoid, the vertex F the Strophoid. This method also easily proves the tangent construction.

Step-by-step description:

1. Let there be two distinct fixed points B and O , both on a given line j . (distance $[B,O]$ will be the radius of the cissoid of Diocle we are about to construct.)
2. Let there be a line k passing O and perpendicular to j .
3. Let there be a circle centered on an arbitrary point C on k , with radius OB .
4. There are two tangents of this circle passing B , let the tangent points be E and F .
5. Let I be the midpoint between E and the center of the circle. Similarly, let J be the midpoint between F and the center of the circle.


6. The locus of I and J (as C moves on k) is the cissoid of Diocles and a line. Also, the locus of E and F is the right strophoid.

Tangent construction for Cissoid and Strophoid: Think of triangle[C,F,B] as a rigid moving body. The point C moves in the direction of vector[O,C], and point B moves in the direction of vector[B,F]. The intersection H (not shown) of normals of line[O,C] and line[B,F] is its center of rotation. J is the point tracing the Cissoid and is also a point on the triangle, thus HJ is normal to the Cissoid. For the Strophoid change the last sentence: Since the tracing point F is a point on the triangle, thus HF is normal to the Strophoid.

In 3D-XploreMath, this construction is shown automatically when Cissoid is chosen from the Plane Curve menu, just after the curve is drawn (or when it is redrawn by choosing Create from the Action menu or typing Command-K). In the Action Menu switch between Cissoid and Strophoid. Hold down the option key to slow the animation, hold down Control to reverse direction, and press the spacebar to pause.


In the animation, the tangent and normal are shown as blue. The critical point that generates the normal line is the intersection of the green lines. One is a vertical drop, the other perpendicular to the red line.

4.4 Pedal and Cardioid




The pedal of a cissoid of Diocles with respect to a point P is the cardioid. If the cissoid's asymptote is the line $y = 1$ and its cusp is at the origin, then P is at $\{0,4\}$. It follows by definition, the negative pedal of a cardioid with respect to a point opposite its cusp is the cissoid of Diocles.

4.5 Negative Pedal and Parabola



The pedal of a parabola with respect to its vertex is the cissoid of Diocles. (and then by definition, the negative pedal of a cissoid of Diocles with respect to its cusp is a parabola.)

4.6 Inversion and Parabola



The inversion of a cissoid of Diocles at cusp is a parabola.

4.7 Roulette of a Parabola

Let there be a fixed parabola. Let there be an equal parabola that rolls on the given parabola in such way that the two parabolas are symmetric to the line of tangency. The vertex of the rolling parabola traces a cissoid of Diocles.

XL.